

# Introducing Mirataz™ (mirtazapine transdermal ointment)

The first and only FDA-approved  
transdermal medication for the  
management of weight loss  
in cats

**"ANOTHER PILL?  
GIVE IT TO THE DOG."**

## Important Safety Information

Mirataz™ (mirtazapine transdermal ointment) is for topical use in cats only under veterinary supervision. Do not use in cats with a known hypersensitivity to mirtazapine or any of the excipients. Do not use in cats treated with monoamine oxidase inhibitors (MAOIs). Not for human use. Keep out of reach of children. Wear gloves when handling/applying, wash hands after and avoid contact between the treated cat and people or other animals for 2 hours following application. Use with caution in cats with hepatic and kidney disease. Cat's food intake should be monitored upon discontinuation. Safety has not been evaluated in cats less than 2 kg, less than six months of age or in breeding, pregnant or lactating cats. The most common adverse reactions observed during clinical trials were application site reactions, behavioral abnormalities (vocalization and hyperactivity) and vomiting. **For product label, including complete safety information, see pages 10-11.**

**Mirataz™**  
(mirtazapine transdermal ointment)

## Identify and manage unintended weight loss in cats

### Weight loss is a clinical sign associated with various underlying diseases in the cat

Asking the right questions can make the difference for the cats in your practice. While there are often subtle changes that the cat owner may or may not have noticed, early detection and action allows the cat to begin eating again, while you diagnose the underlying condition.

#### QUESTIONS TO ASK YOUR CLIENTS

- "Tell me how your cat behaves when he or she eats."*
- "How would you describe your cat's appetite?"*
- "Describe your cat's eating behavior."*
- "How many times does your cat visit the food bowl?"*
- "Describe your typical feeding routine with your cat."*
- "Tell me how your cat's eating habits have changed."*

### Prolonged inadequate nutrition may be more detrimental to the patient than the primary disease process<sup>1</sup>

Therefore, both identifying weight loss and diagnosing the underlying cause are important.

Once changes in eating behavior, body weight, and body condition have been identified and discussed with the cat owner, the tailored diagnostic investigation is initiated, guided by a thorough history and physical exam.


### Some of the more common underlying diseases could be<sup>2-4</sup>:

- ▶ Hyperthyroidism
- ▶ Chronic kidney disease
- ▶ Inflammatory bowel disease
- ▶ Neoplasia
- ▶ Pancreatitis
- ▶ Liver failure

### Important Safety Information

Mirataz™ (mirtazapine transdermal ointment) is for topical use in cats only under veterinary supervision. Do not use in cats with a known hypersensitivity to mirtazapine or any of the excipients or in cats treated with monoamine oxidase inhibitors (MAOIs). Not for human use. Keep out of reach of children. Wear gloves to apply and wash hands after. Avoid contact with treated cat for 2 hours following application. The most common adverse reactions include application site reactions, behavioral abnormalities (vocalization and hyperactivity) and vomiting. **For product label, including complete safety information, see pages 10-11.**

## Mirataz™ (mirtazapine transdermal ointment) manages weight loss in cats



### With Mirataz, veterinarians can confidently recommend an effective topically applied product for cats with unintended weight loss that may be associated with underlying disease

- ✓ Mirataz is the first and only FDA-approved transdermal medication for the management of weight loss in cats<sup>8</sup>
- ✓ In clinical studies, Mirataz resulted in significant weight gain in cats in as little as 14 days following topical application of 2 mg per day<sup>8</sup>
- ✓ Mirataz gives your clients a practical way to manage their cat's weight loss without administration of oral medication and does not rely on the cat to eat to be medicated
- ✓ Due to proprietary Accusorb™ technology, Mirataz achieves measurable plasma concentrations of mirtazapine in cats<sup>9</sup>
- ✓ Mirataz was well tolerated both locally and systemically in clinical studies<sup>8</sup>

# Mirataz™ (mirtazapine transdermal ointment): the first and only FDA-approved transdermal medication for cats


## The active ingredient in Mirataz is mirtazapine

### The pharmacodynamic action of mirtazapine involves antagonism of several receptor sites

Antagonism of presynaptic  $\alpha$ -2 receptors, serotonin receptors (5-HT<sub>2A</sub>, 5-HT<sub>2C</sub>, 5HT<sub>3</sub>), and histamine receptors (H1) by mirtazapine has been demonstrated to result in:

- ▶ Appetite stimulation via interaction with nuclei within the hypothalamus<sup>1</sup>
- ▶ Enhanced release of both serotonin (5-HT) and noradrenaline (NE)<sup>1</sup>
- ▶ Reduced nausea and vomiting in cats<sup>10\*</sup>

\*Vomiting was reported in clinical field studies with the use of Mirataz.



### While not FDA approved, human and compounded versions of mirtazapine have been used off-label but may not be ideal for every cat

- ▶ Human tablets must be split or broken, which results in inaccurate dosing
- ▶ Oral products placed on food will work only if the cat is eating
- ▶ Liquids by mouth may not be any easier than pilling
- ▶ Compounded transdermal mirtazapine has been shown to result in inconsistent and highly variable blood mirtazapine levels<sup>11</sup>


*Mirataz gives your clients an option for one less oral medication for their cats*

### Important Safety Information

Mirataz™ (mirtazapine transdermal ointment) is for topical use in cats only under veterinary supervision. Do not use in cats with a known hypersensitivity to mirtazapine or any of the excipients or in cats treated with monoamine oxidase inhibitors (MAOIs). Not for human use. Keep out of reach of children. Wear gloves to apply and wash hands after. Avoid contact with treated cat for 2 hours following application. The most common adverse reactions include application site reactions, behavioral abnormalities (vocalization and hyperactivity) and vomiting. **For product label, including complete safety information, see pages 10-11.**


# Mirataz achieves measurable blood levels of mirtazapine and effectively results in weight gain in cats

DRUG CONCENTRATION CURVES ( $\pm$ SD) FOR CATS AFTER SINGLE DOSE ORAL ADMINISTRATION AND THE TOPICAL APPLICATION OF MIRATAZ (0.5 MG/KG)<sup>6,9</sup>



### Mirataz demonstrated a 3.9% increase in body weight in cats with unintended weight loss in as little as 14 days<sup>8</sup>

WEIGHT GAIN IN CATS WITH MIRATAZ VS PLACEBO



STUDY DESIGN: Randomized, double-blind, placebo-controlled study (N=177)

### Mirataz was effective in the management of weight loss in cats associated with a wide variety of underlying and concurrent conditions<sup>8</sup>

**Mirataz™**  
(mirtazapine transdermal ointment)

# The safety profile of Mirataz™ (mirtazapine transdermal ointment)

230 cats were enrolled in a field study to assess the clinical safety and effectiveness of Mirataz. Cats enrolled in the study with underlying disease may have received concurrent medications.<sup>6</sup>

## PRE-EXISTING CONDITIONS AND RELEVANT MEDICAL HISTORY OF CATS ENROLLED IN THE STUDY (SAFETY POPULATION)<sup>8</sup>

Description of Condition	Mirataz [n=115 (%)]	Placebo [n=115 (%)]
Any condition	109 (94.8%)	107 (93.0%)
Renal insufficiency	49 (42.6%)	35 (30.4%)
Vomiting	32 (27.8%)	28 (24.3%)
Hyperthyroidism	21 (18.3%)	15 (13.0%)
Dental disease	15 (13.0%)	15 (13.0%)
Heart murmur	15 (13.0%)	13 (11.3%)
Arthritis	15 (13.0%)	11 (9.6%)
Periodontal disorder	12 (10.4%)	15 (13.0%)
Elevated BUN	13 (11.3%)	12 (10.4%)

## CONCOMITANT MEDICATIONS ADMINISTERED (OCCURRING IN >3% OF CATS IN ANY TREATMENT GROUP [EFFECTIVENESS POPULATION])<sup>6</sup>

Concomitant Medication Category	Mirataz [n=115 (%)]	Placebo [n=115 (%)]
Antibiotic	19 (16.5%)	24 (20.9%)
Vitamin/mineral	18 (15.7%)	18 (15.7%)
Parenteral fluids	20 (17.4%)	15 (13.0%)
Anti-thyroid drug	12 (10.4)	9 (7.8%)
Corticosteroid	13 (11.3%)	7 (6.1%)
Vaccine	7 (6.1%)	10 (8.7%)
Anti-hypertensive	8 (7.0%)	9 (7.8%)
Supplement	9 (7.8%)	16 (13.9%)
Anthelmintic or antiparasitic	5 (4.3%)	15 (13.0%)
Antacid	6 (5.2%)	6 (5.2%)
Opioid	6 (5.2%)	8 (7.0%)
Laxative	4 (3.5%)	5 (4.3%)
Otic	3 (2.6%)	2 (1.7%)

- ▶ Beta adrenergic receptor antagonist, anesthetic, antidepressant, and ophthalmic medications were also reported and occurred in <3% of the mirtazapine group<sup>6</sup>

## Mirataz adverse events

The most common adverse reactions observed in the clinical field study included application-site reactions, vocalization, hyperactivity, and vomiting.<sup>8</sup>

- ▶ Application-site reactions resolved upon discontinuation of dosing<sup>8</sup>
- ▶ Of the Mirataz-treated cats that experienced vomiting, 26% had vomiting at the time of enrollment due to underlying conditions<sup>6</sup>

## ADVERSE REACTIONS REPORTED DURING THE CLINICAL FIELD STUDY (OCCURRING IN >3% OF CATS IN THE MIRATAZ GROUP [SAFETY POPULATION])<sup>8</sup>

Description of Adverse Reaction	Mirataz [n=115 (%)]	Placebo [n=115 (%)]
<b>Application Site (ear pinna)</b>		
Erythema	12 (10.4%)	20 (17.4%)
<b>Behavioral</b>		
Vocalization (including crying, mewing)	13 (11.3%)	2 (1.7%)
Hyperactivity (including pacing, restlessness, sleeplessness)	8 (7.0%)	1 (0.9%)
Disoriented state or ataxia	4 (3.5%)	2 (1.7%)
Lethargy (including depressed, sedation, or weakness)	4 (3.5%)	9 (7.8%)
<b>Physical Examination or Observational</b>		
Vomiting	13 (11.3%)	15 (13.0%)
Dehydration	6 (5.2%)	5 (4.3%)
Diarrhea or soft stool	6 (5.2%)	7 (6.1%)
Heart murmur	5 (4.3%)	7 (6.1%)
Inappetence	5 (4.3%)	5 (4.3%)
Renal insufficiency	4 (3.5%)	0
<b>Clinical Pathology</b>		
Hematuria	7 (6.1%)	1 (0.9%)
Elevated BUN (without creatinine)	6 (5.2%)	0
Elevated creatinine and BUN	5 (4.3%)	1 (0.9%)
Hyperphosphatemia	5 (4.3%)	0
Hypokalemia	5 (4.3%)	2 (1.7%)
Pyuria	5 (4.3%)	0

- ▶ Elevated BUN levels were not considered clinically relevant and were likely due to the increased incidence of renal disease (based on clinical pathology and urinalysis) at the time of enrollment in the Mirataz group<sup>8</sup>

## Unintended weight loss can be stressful

Pets bring a great amount of joy to our lives

In fact, **66%** of pet owners believe their pet helps lower stress.<sup>12</sup>


If their cat is experiencing unintended weight loss associated with various underlying conditions, that news alone can be worrisome.

Ensuring at-home treatment plans addressing both the medical needs of the cat and protecting the human-animal bond can be challenging.


With Mirataz™ (mirtazapine transdermal ointment), you can confidently recommend an effective topically applied product for cats with unintended weight loss that may be associated with underlying disease.

## Dosing and administration


Administer topically by applying a 1.5-inch ribbon of ointment (approximately 2 mg/cat) on the inner pinna of the cat's ear once daily for 14 days (see diagrams below)



**Step 1:**  
Wear disposable gloves.  
Twist cap on tube counterclockwise to open.



**Step 2:**  
Apply even pressure on tube and squeeze a 1.5-inch line of ointment onto your gloved finger using the measured line on the carton or in this package insert.



**Step 3:**  
Using your gloved finger, gently rub ribbon of ointment on inside pinna of the cat's ear spreading it evenly over the surface. Dispose of used gloves after each application. If contact with your skin occurs wash thoroughly with soap and warm water.

### Important Safety Information

Mirataz™ (mirtazapine transdermal ointment) is for topical use in cats only under veterinary supervision. Do not use in cats with a known hypersensitivity to mirtazapine or any of the excipients or in cats treated with monoamine oxidase inhibitors (MAOIs). Not for human use. Keep out of reach of children. Wear gloves to apply and wash hands after. Avoid contact with treated cat for 2 hours following application. The most common adverse reactions include application site reactions, behavioral abnormalities (vocalization and hyperactivity) and vomiting. **For product label, including complete safety information, see pages 10-11.**

**Mirataz™**  
(mirtazapine transdermal ointment)

# Mirataz™ (mirtazapine transdermal ointment)

Each 1 g of Mirataz™ contains 20 mg mirtazapine (2%). Each 5 g tube contains 100 mg (0.1 g) of mirtazapine.

For topical application in cats only. Not for oral or ophthalmic use.


## CAUTION:

Federal law (USA) restricts this drug to use by or on the order of a licensed veterinarian.

## DESCRIPTION:

Mirataz™ (mirtazapine transdermal ointment) is a white to off-white ointment containing 2% (w/w) of mirtazapine suitable for transdermal (topical) administration. Mirataz™ contains the following inactive ingredients: Polyethylene Glycol (PEG) 400, PEG 3350, Diethylene Glycol Monoethyl Ether, PEG-8 Caprylic/Capric Glycerides, Oleyl Alcohol, Butylated Hydroxytoluene, Dimethicone, and Dry Flo TS.

The structural formula of mirtazapine is:



Molecular Formula: C<sub>17</sub>H<sub>19</sub>N<sub>3</sub>

Molecular Weight: 265.35

## INDICATION:

Mirataz™ is indicated for the management of weight loss in cats.


## DOSAGE AND ADMINISTRATION:

Administer topically by applying a 1.5-inch ribbon of ointment (approximately 2 mg/cat) on the inner pinna of the cat's ear once daily for 14 days (see Diagrams below).

Wear disposable gloves when applying Mirataz™. Dispose of used gloves after each application.


Alternate the daily application of Mirataz™ between the left and right inner pinna of the ears. Do not administer into the external ear canal. If desired, the inner pinna of the cat's ear may be cleaned by wiping with a dry tissue or cloth immediately prior to the next scheduled dose. If a dose is missed, apply Mirataz™ the following day and resume daily dosing.

To demonstrate the method of administering the dose, the veterinarian or trained personnel at the clinic should apply the first dose in the presence of the owner.




This ruler measures 1.5 inches. Use this ruler to measure the 1.5 inch ribbon of ointment to be applied.


To apply Mirataz™:



**Step 1: Wear disposable gloves. Twist cap on tube counterclockwise to open.**



**Step 2: Apply even pressure on tube and squeeze a 1.5-inch line of ointment onto your gloved finger using the measured line on the carton or in this package insert.**



**Step 3: Using your gloved finger, gently rub ribbon of ointment on inside pinna of the cat's ear spreading it evenly over the surface. Dispose of used gloves after each application. If contact with your skin occurs wash thoroughly with soap and warm water.**

## CONTRAINDICATIONS:

Mirataz™ is contraindicated in cats with a known hypersensitivity to mirtazapine or to any of the excipients.

Mirataz™ should not be given in combination, or within 14 days before or after treatment with a monoamine oxidase inhibitor (MAOI) [e.g. selegiline hydrochloride (L-deprenyl), amitraz], as there may be an increased risk of serotonin syndrome.

## HUMAN WARNINGS:

Not for human use. Keep out of reach of children.

**Wear disposable gloves when handling or applying Mirataz™ to prevent accidental topical exposure.** After application, dispose of used gloves and wash hands with soap and water. After application, care should be taken that people or other animals in the household do not come in contact with the treated cat for 2 hours because mirtazapine can be absorbed transdermally and orally. However, negligible residues are present at the application site and the body of the cat at 2 hours after dosing.

In case of accidental skin exposure, wash thoroughly with soap and warm water. In case of accidental eye exposure, flush eyes with water. If skin or eye irritation occurs seek medical attention.

In case of accidental ingestion, or if skin or eye irritation occurs, seek medical attention.

## PRECAUTIONS:

Do not administer orally or to the eye.

Use with caution in cats with hepatic disease. Mirtazapine may cause elevated serum liver enzymes (See Animal Safety).

Use with caution in cats with kidney disease. Kidney disease may cause reduced clearance of mirtazapine which may result in higher drug exposure.

Upon discontinuation of Mirataz™, it is important to monitor the cat's food intake. Food intake may lessen after discontinuation of mirtazapine transdermal ointment. If food intake diminishes dramatically (>75%) for several days, or if the cat stops eating for more than 48 hours, reevaluate the cat.

Mirataz™ has not been evaluated in cats < 2 kg or less than 6 months of age. The safe use of Mirataz™ has not been evaluated in cats that are intended for breeding, pregnant, or lactating cats.

## ADVERSE REACTIONS:

In a randomized, double-masked, vehicle-controlled field study to assess the effectiveness and safety of mirtazapine for the management of weight loss in cats, 115 cats treated with Mirataz™ and 115 cats treated with vehicle control were evaluated for safety. The vehicle control was an ointment containing the same inert ingredients as Mirataz™ without mirtazapine. The most common adverse reactions included application site reactions, behavioral abnormalities (vocalization and hyperactivity), and vomiting. The adverse reactions observed in the study and number of cats experiencing each adverse reaction is summarized in Table 1 below.

**Table 1. Adverse Reactions Reported During the Field Study**

Adverse Reaction	Mirataz™ N=115 (%)	Vehicle Control N=115 (%)
<b>Application site (Ear pinna)</b>		
Erythema	12 (10.4%)	20 (17.4%)
Crust/Scab	3 (2.6%)	6 (5.2%)
Residue	3 (2.6%)	8 (7.0%)
Scaling/Dryness	3 (2.6%)	3 (2.6%)
Dermatitis or irritation	1 (0.9%)	9 (7.8%)
Alopecia	1 (0.9%)	2 (1.7%)
Pruritus	1 (0.9%)	4 (3.5%)
<b>Behavioral</b>		
Vocalization	13 (11.3%)	2 (1.7%)
Hyperactivity	8 (7.0%)	1 (0.9%)
Disoriented state or ataxia	4 (3.5%)	2 (1.7%)
Lethargy/weakness	4 (3.5%)	9 (7.8%)
Attention Seeking	3 (2.6%)	0
Aggression	2 (1.7%)	0
<b>Physical Examination or Observational</b>		
Vomiting	13 (11.3%)	15 (13.0%)
Dehydration	6 (5.2%)	5 (4.3%)
Diarrhea or soft stool	6 (5.2%)	7 (6.1%)
Heart murmur	5 (4.3%)	7 (6.1%)
Inappetence	5 (4.3%)	5 (4.3%)
Renal insufficiency*	4 (3.5%)	0
Ear infection	3 (2.6%)	0
Urinary tract infection	3 (2.6%)	0
<b>Clinical Pathology</b>		
Hematuria	7 (6.1%)	1 (0.9%)
Elevated BUN (without creatinine)**	6 (5.2%)	0
Elevated creatinine and BUN	5 (4.3%)	1 (0.9%)
Hyperphosphatemia	5 (4.3%)	0
Hypokalemia	5 (4.3%)	2 (1.7%)
Pyuria	5 (4.3%)	0
Anemia	3 (2.6%)	8 (7.0%)
Low urine specific gravity	3 (2.6%)	1 (0.9%)
Monocytosis	3 (2.6%)	2 (1.7%)
Neutrophilia	3 (2.6%)	2 (1.7%)

\* One cat with renal insufficiency was reported with a serious adverse reaction of acute renal failure, hematuria, and pyuria at the Week 2 visit. The cat was enrolled with a history of chronic kidney disease. Euthanasia was elected and necropsy revealed hypertrophic cardiomyopathy, bilateral parathyroid hyperplasia, and mild to moderate renal disease.

\*\* At Week 2, blood urea nitrogen (BUN) values were significantly higher in the Mirataz™ group compared to the vehicle control group (p<0.10). The BUN in the Mirataz™ group was 43.60 mg/dL (reference range 16-37 mg/dL) compared to 36.05 mg/dL in the vehicle control group.

Post study, follow-up was done in 199 cats (103 in the Mirataz™ (mirtazapine transdermal ointment) group and 96 in the vehicle control group). Following cessation of Mirataz™, four cats were reported as being less social or less restless, one cat was reported as more active, and one cat was reported with increased hissing and urinating out of the litter box.

To report suspected adverse events, for technical assistance or to obtain a copy of the SDS, contact Kindred Biosciences, Inc. at 888-608-2542.

For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at <http://www.fda.gov/AnimalVeterinary/SafetyHealth>

## INFORMATION FOR CAT OWNERS:

Upon discontinuation of Mirataz™, it is important to monitor your cat's food intake. Your cat's food intake may lessen after discontinuation of mirtazapine transdermal ointment. If food intake diminishes dramatically (>75%) for several days or if your cat stops eating for more than 48 hours, contact your veterinarian.

## CLINICAL PHARMACOLOGY:

### Mechanism of Action:

The exact mechanism by which mirtazapine induces weight gain has not been clearly elucidated but appears to be multifactorial.<sup>1</sup> Mirtazapine is an  $\alpha$ 2-adrenergic receptor antagonist nor-adrenergic and serotonergic antidepressant drug. Mirtazapine is known to be a potent antagonist of 5-HT2 and 5-HT3 serotonin receptors in the central nervous system (CNS), and a potent inhibitor of histamine H1 receptors. Because mirtazapine blocks 5-HT2 and 5-HT3 receptors, only 5-HT1A-mediated serotonergic transmission is enhanced. Inhibition of 5-HT2 receptors may account for the orexigenic effects of mirtazapine. Another hypothesis is that mirtazapine induced weight gain may be secondary to changes in leptin and the tumor necrosis factor  $\alpha$ (TNF- $\alpha$ ) cytokine system.<sup>2</sup> A study by Fernstrom (1995) demonstrated a reduction of the basal metabolic rate in patients treated with antidepressants in general.<sup>3</sup>

### Pharmacokinetics:

In a crossover study in eight cats to determine the relative bioavailability of oral and transdermal 2% mirtazapine, the mean half-life (26.8 hours) with topical administration was over 2X longer than the mean half-life (10.1 hours) with oral administration.

The doses used in the target animal safety study were higher (2.8 to 5.4 mg) than the label dose. Based on dose proportionality in AUC<sub>0-24hr</sub> and C<sub>max</sub> observed in this study, these pharmacokinetic parameters were extrapolated for the 2 mg/cat label dose administered once per day for 35 days (see Table 2).

Steady state was achieved within 14 days. The median accumulation between first and 35<sup>th</sup> dose was 3.71X (based on AUC ratio) and 3.90X (based on C<sub>max</sub> ratio).

**Table 2. Plasma pharmacokinetic parameters at steady state after 2 mg/cat dose of mirtazapine 2.0% transdermal ointment in healthy cats**

Parameter	Unit	Mean (SD)
C <sub>max</sub>	ng/mL	32.1 (19.9)
T <sub>max</sub>	hr	6 (2-6) <sup>a</sup>
AUC <sub>0-24hr</sub>	hr*ng/mL	410.3 (213.5)
Half-life <sup>b</sup>	hr	11.2 (2.98)

C<sub>max</sub> = extrapolated maximum plasma concentration

<sup>a</sup>T<sub>max</sub> = time to maximum plasma concentration (reported as median and range)

AUC<sub>0-24hr</sub> = extrapolated area under the plasma time vs. concentration curve

<sup>b</sup>The half-life value is reflective of both topical and oral exposure. In another study where cats wore Elizabethan collar to restrict access to their ears and consequent oral exposure, a longer half-life (Mean= 20.7 hr) was observed

## EFFECTIVENESS:

The effectiveness of Mirataz™ (mirtazapine transdermal ointment) was demonstrated in a randomized, double-masked, vehicle-controlled, multi-site field study involving client-owned cats of various breeds.

Enrolled cats were  $\geq$  1 year of age and had existing documented medical history of  $\geq$  5% weight loss deemed clinically significant. The most common pre-existing conditions included renal insufficiency, vomiting, and hyperthyroidism. Some cats had more than one pre-existing condition. Cats were randomized to treatment groups in a 1:1 ratio of Mirataz™ to vehicle control. A total of 230 cats were enrolled and received either Mirataz™ (115 cats) or a vehicle control (115 cats) containing the same inert ingredients without mirtazapine. The cats were 2.8-24.6 years of age and weighed 2.1-9.2 kg. The dosage was a 1.5-inch ribbon (approximately 2 mg/cat) mirtazapine or vehicle ointment administered topically to the inner pinna of the cat's ear.

A total of 177 cats were determined to be eligible for the effectiveness analysis; 83 cats were in the Mirataz™ group and 94 cats were in the vehicle control group. The primary effectiveness endpoint was the mean percent change in body weight from Day 1 to the Week 2 Visit.

At Week 2, the mean percent increase in body weight from Day 1 was 3.94% in the mirtazapine group and 0.41% in the vehicle control group. The difference between the two groups was significant (p<0.0001) based on a two-sample t-test assuming equal variances. A 95% confidence interval on the mean percent change in body weight for the Mirataz™ group is (2.77, 5.11), demonstrating that the mean percent change is statistically different from and greater than 0.

## ANIMAL SAFETY:

The margin of safety of mirtazapine was evaluated in one laboratory study, a comprehensive review of six pilot studies (five laboratory and one clinical) utilizing the final market formulation, and one laboratory study that was not final market formulation.

## Laboratory Safety Study:

In a 6-week laboratory safety study, 48 healthy cats aged 7-10 months were dosed topically with mirtazapine once daily at 0 mg/kg (vehicle control), 1.1 mg/kg (1.4 to 2.7X), 3.2 mg/kg (4.3-7.5X), and 5.3 mg/kg (7.1-12.2X) body weight. Four cats/sex/group in the 1.1 and 3.2 mg/kg groups were

dosed topically to the inner pinna of the ear, alternating between right and left ears. Eight cats/sex/group in the 0 and 5.3 mg/kg groups were dosed topically to the inner pinna of the ear, splitting the dose between both ears. Four cats/sex/group in the 0 and 5.3 mg/kg groups were maintained and monitored during a 4-week recovery period.

## 6 Week Dosing Period

Application of mirtazapine and vehicle control was associated with ear flicking, head shaking, pulling away/flinching and infrequently with struggling/fractioning behavior, and hypersalivation. Inner and outer pinna erythema, flaking, alopecia, and thickening were observed in all cats in all groups. Erythema, crusting, alopecia, and scabbing of the skin, mostly around the head and neck, was frequently observed in all groups and occasionally affected the tail, tarsi or carpi, likely due to spread of the ointment to these areas by self-grooming.

Mirtazapine administration resulted in increased vocalization, hyperactivity, attention-seeking behaviors, and tremors in all mirtazapine dose groups. Frank blood in the stool was infrequently observed in the vehicle control, 3.2, and 5.3 mg/kg groups. Polyuria was observed in all groups. Polydipsia was observed in one cat in the 5.3 mg/kg group. Three cats (one from each mirtazapine dose group) were isosthenuric. Eight cats developed cystitis with or without urethral obstruction throughout the study in all groups. One cat each from the vehicle control and 3.2 mg/kg group were euthanized early on Day 35 due to urethral obstruction.

Two cats from the 1.1 mg/kg group had either ventricular premature contractions (VPC) or tall R waves, and one cat from the 5.3 mg/kg group had both VPC and a right axis deviation.

Eosinophilia was noted sporadically in the vehicle control, 1.1, and 5.3 mg/kg groups. Mild elevations in ALT values were noted sporadically in vehicle control, 3.2, and 5.3 mg/kg groups. On Day 15, one cat in the 3.2 mg/kg group demonstrated a marked ALT elevation of 3397 U/L, with concurrent elevations in AST and GGT. By Day 42, the ALT declined to 109 U/L and the AST and GGT returned to within normal limits.

Gross pathology findings, confirmed with histopathology, were hyperplastic dermatitis (alopecia, hyperkeratosis, thickening, and ceruminous gland secretion) of the pinnae in all cats and findings consistent with cystitis (mucosal urinary bladder hemorrhage, mottled-dark red appearance, and irregular contour) in four male cats (two vehicle control, one 3.2 mg/kg and one 5.3 mg/kg). Additional histopathologic findings included pyelonephritis (two vehicle control and one 5.3 mg/kg), nephrocalcinosis (three vehicle control, one 1.1 mg/kg, three 3.2 mg/kg, and one 5.3 mg/kg), necrosis of the kidneys (one vehicle control and one 3.2 mg/kg), unilateral hypoplasia of the thyroid gland (two 5.3 mg/kg), and unilateral hypertrophy of the thyroid gland (one 5.3 mg/kg).

## 4 Week Recovery Period

Following a 4-week recovery period, ALT elevations resolved. Polyuria was reduced and only occurred in two cats in the 5.3 mg/kg group. Pinnal lesions (erythema and flaking) completely resolved in the vehicle control and improved in the 5.3 mg/kg groups. Ear thickening improved in both groups.

## Pilot Safety Studies:

In six pilot studies (five laboratory and one clinical study) utilizing the final market formulation of Mirataz™, and one laboratory study that utilized non-final market formulation, a total of 76 cats were administered mirtazapine. Five studies administered mirtazapine ointment topically (0.5-5.3 mg/kg), one study administered mirtazapine topically and orally (0.5 mg/kg), and one study administered mirtazapine orally (10.6 mg/cat).

The most common observations were ear pinnae reactions (erythema with or without blood and flaking), mild behavioral observations (vocalization and tremors), vomiting, and diarrhea.

One study reported six cats with blood in the stool. Two studies reported one cat each showing aggression. One study reported polyuria in one cat; another study reported stranguria and possible urinary tract infection in one cat. Two studies reported cardiac abnormalities, including sinus tachycardia not present at baseline, development of a grade 3/6 heart murmur, tall QRS complexes, and left or right axis deviation. One study reported one cat with ataxia. In the studies that administered mirtazapine orally, salivation and lip licking were frequently observed.

In the terminal study administering 5.3 mg/kg mirtazapine topically, histopathology showed chronic hyperplastic dermatitis at the application site.

## STORAGE:

Store below 25°C (77°F). Multi-use tube. Discard within 30 days of first use.

## HOW SUPPLIED:

Mirataz™ is supplied in a 5 gram aluminum tube.

## REFERENCES:

- Laimer, M., Kramer-Reinstadler, K., Rauchenzauner, M., Lechner-Schoner, T., Strauss, R., Engl, J., & Ebenbichler, C. F. (2006). Effect of mirtazapine treatment on body composition and metabolism. *The Journal of clinical psychiatry*, 67(3), 421-424.
- Kraus, T., Haack, M., Schuld, A., Hinze-Selch, D., Koethe, D., & Pollmächer, T. (2002). Body weight, the tumor necrosis factor system, and leptin production during treatment with mirtazapine or venlafaxine. *Pharmacopsychiatry*, 35(06), 220-225.
- Fernstrom, M. H. (1995). Drugs that cause weight gain. *Obesity research*, 3(S4), 435S-439S.

## NADA 141-481, Approved by FDA

NDC 86078-686-01

REG-007 Rev. 11/17

## MANUFACTURED FOR:

Kindred Biosciences, Inc.  
1555 Bayshore Highway, suite 200  
Burlingame, CA 94010


Made in USA.

Mirataz™ is a trademark of Kindred Biosciences, Inc.  
©2018 Kindred Biosciences, Inc. All rights reserved.

KindredBio

# Manage weight loss in cats with Mirataz™ (mirtazapine transdermal ointment)

- ✓ Mirataz is the first and only FDA-approved transdermal medication for the management of weight loss in cats
- ✓ In clinical studies, Mirataz resulted in significant weight gain in cats in as little as 14 days following topical application of 2 mg per day<sup>8</sup>
- ✓ Mirataz gives your clients a practical way to manage their cat's weight loss without administration of oral medication and does not rely on the cat to eat to be medicated
- ✓ Due to proprietary Accusorb™ technology, Mirataz achieves measurable plasma concentrations of mirtazapine in cats<sup>9</sup>
- ✓ Mirataz was well tolerated both locally and systemically in clinical studies<sup>8</sup>



**"ANOTHER PILL?  
GIVE IT TO THE DOG."**

For more information, contact your KindredBio Sales Specialist at 1-888-608-2542, your preferred Distributor Sales Representative, or go to [kindredbio.com/Mirataz](https://www.kindredbio.com/Mirataz)

## Important Safety Information

Mirataz™ (mirtazapine transdermal ointment) is for topical use in cats only under veterinary supervision. Do not use in cats with a known hypersensitivity to mirtazapine or any of the excipients. Do not use in cats treated with monoamine oxidase inhibitors (MAOIs). Not for human use. Keep out of reach of children. Wear gloves when handling/applying, wash hands after and avoid contact between the treated cat and people or other animals for 2 hours following application. Use with caution in cats with hepatic and kidney disease. Cat's food intake should be monitored upon discontinuation. Safety has not been evaluated in cats less than 2 kg, less than six months of age or in breeding, pregnant or lactating cats. The most common adverse reactions observed during clinical trials were application site reactions, behavioral abnormalities (vocalization and hyperactivity) and vomiting. **For product label, including complete safety information, see pages 10-11.**

**References:** 1. Agnew W, Korman R. Pharmacological appetite stimulation: rational choices in the inappetent cat. *J Feline Med Surg.* 2014;16(9):749-756. 2. Pittari J, Rodan I, Beekman G, et al. American association of feline practitioners. Senior care guidelines. *J Feline Med Surg.* 2009;11(9):763-778. 3. Laflamme DP. Nutrition for aging cats and dogs and the importance of body condition. *Vet Clin North Am Small Anim Pract.* 2005;35(3):713-742. 4. Baez JL, Michel KE, Sorenmo K, et al. A prospective investigation of the prevalence and prognostic significance of weight loss and changes in body condition in feline cancer patients. *J Feline Med Surg.* 2007;9(5):411-417. 5. 2017 Mirataz Pricing Research, Kynotec, September 2017 (n=204 U.S. small animal veterinarians). 6. Data on file. Kindred Biosciences, Inc. 7. American Pet Products Association (APPA), 2017-2018 National Pet Owners Survey (n=22,202 respondents). 8. Mirataz™ (mirtazapine transdermal ointment) [package insert], Kindred Biosciences, Inc. (Burlingame, CA). Rev. 5/2018. 9. Buhles W, Quimby JM, Labelle D, et al. Single and multiple dose pharmacokinetics of a novel transdermal ointment in cats. *J Vet Pharmacol Ther.* In press 2018. 10. Quimby JM, Lunn KF. Mirtazapine as an appetite stimulant and anti-emetic in cats with chronic kidney disease: a masked placebo-controlled crossover clinical trial. *Vet J.* 2013;197(3):651-655. 11. Benson KK, Zajic LB, Morgan PK, et al. Drug exposure and clinical effect of transdermal mirtazapine in healthy young cats: a pilot study. *J Feline Med Surg.* 2017;19(10):998-1006. 12. American Pet Products Association. The 2017-2018 APPA National Pet Owner's Survey video. American Pet Products Association website. <https://vimeo.com/orangetreeps/review/214716969/38f274fe6d>. Accessed on December 5, 2017.



Mirataz™ and Accusorb™ are trademarks of KindredBio  
©2018 Kindred Biosciences, Inc., Burlingame, CA 94010. All rights reserved  
US-MAZ-1800011 5/4/18

**Mirataz™**  
(mirtazapine transdermal ointment)